

Research



**Cite this article:** Mathon L *et al.* 2022 Cross-ocean patterns and processes in fish biodiversity on coral reefs through the lens of eDNA metabarcoding. *Proc. R. Soc. B* **289**: 20220162.  
<https://doi.org/10.1098/rspb.2022.0162>

Received: 27 January 2022  
 Accepted: 24 March 2022

**Subject Category:**

Ecology

**Subject Areas:**

biotechnology, ecology

**Keywords:**

eDNA metabarcoding, coral reef fish, biogeographic patterns, visual census

**Author for correspondence:**

Laetitia Mathon  
 e-mail: [laetitia.mathon@gmail.com](mailto:laetitia.mathon@gmail.com)

†These authors contributed equally as first author to this work.

‡These authors contributed equally as senior author to this work.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5933156>.

# Cross-ocean patterns and processes in fish biodiversity on coral reefs through the lens of eDNA metabarcoding

Laetitia Mathon<sup>1,2,†</sup>, Virginie Marques<sup>1,3,†</sup>, David Mouillot<sup>3,4</sup>, Camille Albouy<sup>5</sup>, Marco Andrello<sup>3,17</sup>, Florian Baletaud<sup>2,3,6</sup>, Giomar H. Borrero-Pérez<sup>7</sup>, Tony Dejean<sup>8</sup>, Graham J. Edgar<sup>9</sup>, Jonathan Grondin<sup>8</sup>, Pierre-Edouard Guerin<sup>1</sup>, Régis Hocdé<sup>3</sup>, Jean-Baptiste Juhel<sup>3</sup>, Kadarusman<sup>10</sup>, Eva Maire<sup>3,11</sup>, Gael Mariani<sup>3</sup>, Matthew McLean<sup>12</sup>, Andrea Polanco F.<sup>7</sup>, Laurent Pouyau<sup>13</sup>, Rick D. Stuart-Smith<sup>9</sup>, Hagi Yulia Sugeha<sup>14</sup>, Alice Valentini<sup>8</sup>, Laurent Vigliola<sup>2</sup>, Indra B. Vimono<sup>14</sup>, Loïc Pellissier<sup>15,16,‡</sup> and Stéphanie Manel<sup>1,‡</sup>

- <sup>1</sup>CEFE, Univ. Montpellier, CNRS, EPHE-PSL University, IRD, Montpellier, France
- <sup>2</sup>ENTROPIE, Institut de Recherche pour le Développement (IRD), Univ. Réunion, UNC, CNRS, Q1 IFREMER, Nouméa, New Caledonia, France
- <sup>3</sup>MARBEC, Univ Montpellier, CNRS, IFREMER, IRD, Montpellier, France
- <sup>4</sup>Institut Universitaire de France, France
- <sup>5</sup>DECOD (Ecosystem Dynamics and Sustainability), IFREMER, INRAE, Institut Agro - Agrocampus Ouest, Nantes, France
- <sup>6</sup>SOPRONER, groupe GINGER, 98000 Noumea, New Caledonia, France
- <sup>7</sup>Programa de Biodiversidad y Ecosistemas Marinos, Museo de Historia Natural Marina de Colombia (MHNMC), Instituto de Investigaciones Marinas y Costeras- INVEMAR, Santa Marta, Colombia
- <sup>8</sup>SPYGEN, Le Bourget-du-Lac, France
- <sup>9</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania, Australia
- <sup>10</sup>Politeknik Kelautan dan Perikanan Sorong, KKD BP Sumberdaya Genetik, Konservasi dan Domestikasi, Papua Barat, Indonesia
- <sup>11</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK
- <sup>12</sup>Department of Biology, Dalhousie University, Halifax NSB3H4R2, Canada
- <sup>13</sup>ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France
- <sup>14</sup>Research Center for Oceanography, National Research and Innovation Agency, Jl. Pasir Putih 1, Ancol Timur, Jakarta Utara 14430, Indonesia
- <sup>15</sup>Landscape Ecology, Institute of Terrestrial Ecosystems, Department of Environmental Systems Science, ETH Zürich, Zürich, Switzerland
- <sup>16</sup>Unit of Land Change Science, Swiss Federal Research Institute WSL, Birmensdorf, Switzerland
- <sup>17</sup>Institute for the study of Anthropic Impacts and Sustainability in the marine environment, National Research Council (CNR-IAS), Rome, Italy

LM, 0000-0001-8147-8177; DM, 0000-0003-0402-2605; MA, 0000-0001-7590-2736; AV, 0000-0001-5829-5479; LP, 0000-0002-2289-8259; SM, 0000-0001-8902-6052

Increasing speed and magnitude of global change threaten the world's biodiversity and particularly coral reef fishes. A better understanding of large-scale patterns and processes on coral reefs is essential to prevent fish biodiversity decline but it requires new monitoring approaches. Here, we use environmental DNA metabarcoding to reconstruct well-known patterns of fish biodiversity on coral reefs and uncover hidden patterns on these highly diverse and threatened ecosystems. We analysed 226 environmental DNA (eDNA) seawater samples from 100 stations in five tropical regions (Caribbean, Central and Southwest Pacific, Coral Triangle and Western Indian Ocean) and compared those to 2047 underwater visual censuses from the Reef Life Survey in 1224 stations. Environmental DNA reveals a higher (16%) fish biodiversity, with 2650 taxa, and 25% more families than underwater visual surveys. By identifying more pelagic, reef-associated and crypto-benthic species, eDNA offers a fresh view on assembly rules across spatial scales. Nevertheless, the reef life survey identified more species than eDNA in 47 shared families, which can be due to incomplete

sequence assignment, possibly combined with incomplete detection in the environment, for some species. Combining eDNA metabarcoding and extensive visual census offers novel insights on the spatial organization of the richest marine ecosystems.

## 1. Introduction

Coral reefs host the highest fish diversity on earth despite covering less than 0.1% of the ocean's surface [1,2]. They are also severely threatened [3], with near-future outlooks predominantly pessimistic [4]. Data syntheses over decades of surveys estimate the total number of coral reef fishes to be 2400 to 8000 species [5,6], distributed among approximately 100 families [7]. Typically, coral reef biodiversity displays clear spatial patterns, including longitudinal and latitudinal gradients outwards the Indo-Australian Archipelago [8,9], also known as the 'Coral Triangle', hosting the world's highest level of marine biodiversity [10]. The exceptional biodiversity in the Coral Triangle has recently been suggested to strongly relate to higher diversity among fish families that feed on plankton [11]. Other trophic groups are also very important on coral reefs but are often undetected because they are transient or hidden [12,13]. Intriguingly, the proportions of fish species among families are shown to be strongly conserved across the Indo-Pacific [8]. The spatial patterns of coral reef fishes are also marked by strong variations in taxonomic composition (species turnover or  $\beta$  diversity), often due to isolation [14]. Many species on coral reefs are geographically localized, but can sometimes be locally abundant, while others are widespread [15].

Coral reef fishes have evolved in a physically complex environment and present a wide range of forms and functions [16]. Small cryptic species, hereafter called crypto-benthic, that live inside the reef structure, can be very difficult to sample or survey using non-destructive methods [17], yet represent half of the fish diversity on coral reefs [13]. Even though fishes are among the best-studied taxa inhabiting coral reefs [18], our knowledge of their biodiversity is only partial [19], the taxonomy is complex, uncertain for many species [5], and countless species remain undescribed.

Environmental DNA (eDNA) metabarcoding, a method retrieving and analysing DNA naturally released by organisms in their environment [20], provides an opportunity to not only better understand classical biodiversity patterns, but also uncover novel ones hidden by our incomplete taxonomic and biogeographic coverage [21]. Environmental DNA is particularly powerful in aquatic ecosystems [22] and is now well established for marine microorganisms [23,24]. By contrast, its potential to provide an integrated biodiversity assessment of macroorganisms, including vertebrates of all trophic levels (from crypto-benthic to large pelagic fish species), is only shown at local [25] and regional [26–30] scales, but not yet at spatial scales including more than one biogeographic region or multiple ocean basins.

Here, we investigate how a cross-ocean basin snapshot of eDNA sampling could describe the distribution of fish biodiversity on coral reefs, reveal unknown patterns and challenge well-established assembly rules. From 226 eDNA seawater samples (2712 PCR replicates) collected in 100 stations at 26 sites covering five tropical regions (Southeast Polynesia,


Tropical Northwestern Atlantic, Tropical Southwestern Pacific, Western Indian Ocean and Western Coral Triangle) across the Indian, Pacific and Atlantic Oceans (electronic supplementary material, figures S1 and S2), we produced a final dataset of 189 350 273 mitochondrial 12S rRNA gene sequence reads (see Methods), clustered into 2023 molecular operational taxonomic units (MOTUs) and assigned to Actinopterygii (bony fishes) and Chondrichthyes (cartilaginous fishes) taxa (electronic supplementary material, tables S1 and S2). We then compared fish biodiversity patterns obtained from eDNA to those observed from 2047 standardized visual surveys of reef fishes in 1224 stations at 219 sites within 24 tropical regions [31].

## 2. Results

### (a) Global estimates of fish biodiversity on coral reefs

We estimated total fish diversity on coral reefs using the asymptote of a multi-model accumulation curve for both eDNA MOTUs [32] and visual census species (see Methods). The asymptote estimated from 100 eDNA stations distributed in five regions sampled over a 28-month period reaches 2650 MOTUs (figure 1a). This detectable fish MOTU diversity, including also MOTUs unassigned at the species level, is 16% higher than the estimate from visual census data, which reaches an asymptote at 2268 fish species from 2047 tropical transects surveyed during 13 years (figure 1b). The asymptotic estimation of family richness obtained with eDNA reaches 147 families, 25% more than the asymptotic number of families estimated with visual census data (118 families, figure 1c,d). Among the 71 families shared between both datasets, 24 have a higher number of MOTUs from eDNA survey than species from the visual survey while 47 have more species from visual survey than MOTUs from eDNA survey (figure 1e). Families with more taxa identified using eDNA include those often associated with reef-adjacent habitats such as mangroves or soft sediments like Mugilidae (e.g. *Mugil rubrioculus*), Elopidae and Gerreidae [33] (e.g. *Gerres oyna*), and crypto-benthic species that live hidden in crevices (e.g. Gobiidae) or nocturnal fish species [34] (e.g. Congridae). Families with more taxa with visual census include Acanthuridae, Chaetodontidae, Blenniidae, Labridae, Pomacentridae and Scaridae. Fifty-five families are detected only with eDNA, including Myctophidae, Engraulidae, Atherinidae and Exocoetidae, while 24 families are detected only by the visual census, including Caesionidae, Chaenopsidae, Labrisomidae and Microdesmidae. Environmental DNA estimates a diversity of crypto-benthic species 13% higher than with visual census, and, among many others, includes species such as the elegant firefish (*Nemateleotris decora*), which lives on the outer reef slope between 25 and 70 m (figure 2a). Yet, the difference in fish diversity assessment between the two methods is the strongest for pelagic and wide-ranging species, for which eDNA reveals more than seven times higher richness than with visual census. These species mainly belong to Scombridae (e.g. *Katsuwonus pelamis*), Clupeidae, Carcharhinidae (e.g. *Carcharhinus leucas*, *Sphyrna lewini*) and Belonidae (figure 2b).

MOTU richness per fish family retrieved with eDNA is strongly correlated with fish species richness within families recorded in visual census data (Pearson correlation = 0.84,  $p < 0.001$ ,  $n = 71$ ; figure 1e). Highly diverse families seen on coral reefs are also well represented in eDNA samples, with




**Figure 1.** Estimates of overall fish richness from environmental DNA (eDNA) and visual census. (a) Accumulation curve of MOTUs from eDNA (eDNA MOTUs), (b) accumulation curve of species from the visual census database, (c) accumulation curve of eDNA families and (d) accumulation curve of visual census families. For (a–d), species accumulation model is fitted according to Lomolino method (see methods). (e) Linear regression (black line) between the number of species per family in visual census data and the number of MOTUs per family in eDNA ( $\log(x + 1)$  transformation) over  $n = 77$  families. Each point is a family. Red line is  $x = y$ . (f) percentage of MOTUs assigned to each family at global scale and proportion in each region. (no. = number of) (Online version in colour.)

Gobiidae, Labridae and Pomacentridae containing more than 100 MOTUs each, together representing about 20% of MOTUs (figure 1f; electronic supplementary material, figures S3 and S4). The slope of the log–log relationship between MOTUs richness per family and species richness per family is equal to 0.8 showing that the relationship is not proportional but saturating. The richest fish families contain more MOTUs detected with eDNA than species detected with visual surveys.

### (b) Biogeography of eDNA sequences

The spatial distribution of MOTUs follows clear biogeographic patterns, with a peak in the Coral Triangle and lower values of MOTU richness toward Southeast Polynesia (electronic supplementary material, figure S5). The richest region (West Papua, Indonesia, Western Coral Triangle) contains

approximately 50% of the global pool of fish MOTUs while the poorest region (Fakarava, French Polynesia, Southeast Polynesia) contains only 9% of the global pool (electronic supplementary material, figures S6 and S7, and table S2). Distance-based redundancy analysis (dbRDA) was performed on fish family proportions at each site (i.e. number of MOTUs or species assigned to each family in each site, see Methods) for eDNA and visual surveys with the region and the site MOTU/species richness as explanatory variables, including their interaction (figure 3; electronic supplementary material, table S3). For eDNA, the dbRDA explains up to 42% of variation in family proportions between pairs of sites with region and MOTU/species richness both having significant effects ( $F = 4.1$  and  $5.7$ , respectively,  $p < 0.001$ ), but no significant interaction ( $F = 1.99$ ,  $p > 0.05$ ). The partial dbRDA on eDNA showed a significant effect of region while controlling for MOTU



**Figure 2.** Estimates of overall fish richness from eDNA and visual census across habitat categories. (a) Accumulation curve of crypto-benthic eDNA MOTUs (i) and visual census species (ii). (b) Accumulation curve of pelagic MOTUs (i) and visual census species (ii). (c) Accumulation curve of demersal MOTUs (i) and visual census species (ii). Accumulation model is fitted with a nonlinear Lomolino model (see Methods). (Online version in colour.)


richness ( $F = 2.79$ ,  $p < 0.001$ ). The first axis explains 17.2% of variation in family proportions and separates the Western Coral Triangle from other regions (figure 3*a,b*). The first axis shows a higher proportion of Lutjanidae but lower proportions of Labridae and Gobiidae in sites of the Western Coral Triangle. It also confirms the longitudinal diversity gradient from the Coral Triangle. The second axis explains 11.2% of variation and discriminates the Tropical Northwestern Atlantic from the Western Indian Ocean, due to a higher proportion of Clupeidae and Carangidae in the Atlantic Ocean and a higher proportion of Acanthuridae in the Indian Ocean. The dbRDA performed on visual census data explained greater variation ( $R^2 = 0.5$ ,  $p < 0.001$ ) and the region also had a significant, albeit weaker than for MOTUs, effect on fish family proportions ( $F = 17.7$ ,  $p < 0.01$ ), while species richness and interaction between the two variables also had significant effects ( $F = 6.28$  and  $2$ ,  $p < 0.01$ , respectively). The first axis

explains 41.6% of variance in family proportions and separates the Tropical Northwestern Atlantic from the other regions with a higher proportion of Gobiidae and Serranidae. The second axis explains 5.7% of variance in family proportions and separates the Southeast Polynesia from Indo-Pacific regions, and is mostly driven by the higher proportion of Pomacentridae in the Indo-Pacific (figure 3*c,d*).

### (c) Global patterns of fish turnover and rarity

Our eDNA survey shows that a majority of MOTUs are geographically restricted, with 85% of the MOTUs detected in only one region (figure 4*a*), and 35% in only one site (electronic supplementary material, figure S8). Geographic restriction is one aspect of species rarity but is shown to play a primary role in determining extinction risk while local abundance and habitat specialization have secondary





**Figure 3.** Partial dbRDA of MOTU proportions of each family in each site. (a) dbRDA on eDNA dataset, with 133 families in 26 sites ( $R^2 = 0.21$ ,  $F = 3.11$ ,  $p = 0.001$ ). (b) Families with scores greater than 95% of scores distribution on each axis for eDNA. (c) dbRDA on a subset of visual census dataset to select only the sites in the same regions as in the eDNA dataset, with 76 families in 68 sites ( $R^2 = 0.5$ ,  $F = 15.8$ ,  $p = 0.001$ ). (d) Families with scores greater than 95% of scores distribution on each axis for visual census. Axis labels indicate the percentage of variance explained by the 2 first dbRDA dimensions (CAP1 and CAP2). (Online version in colour.)

roles [35]. We hierarchically partitioned the global MOTU diversity ( $\gamma_{\text{global}}$ ) into additive diversity components (i.e. dissimilarity) due to difference between regions ( $\beta_{\text{inter-region}}$ ), mean difference between sites within regions ( $\bar{\beta}_{\text{inter}}$ ), mean difference between stations within sites ( $\bar{\beta}_{\text{inter-station}}$ ) and mean station diversity ( $\bar{\alpha}_{\text{station}}$ ) [36]. As a consequence of the geographic restriction of most MOTUs to one region, the total fish MOTU ( $\gamma$ ) diversity is mainly due to inter-region  $\beta$ -diversity (approx. 74%) followed by inter-site (14.8%) and inter-station (5.9%)  $\beta$ -diversity (figure 4b). The same partitioning using different site delineations (10 and 20 km) provides similar results (electronic supplementary material, table S4). Diversity partitioning of crypto-benthic fish MOTUs only or pelagic fish MOTUs only reveals similar patterns (electronic supplementary material, table S5). The partitioning diversity of species detected by visual census also revealed similar patterns but with a stronger effect of  $\beta_{\text{inter-region}}$  (84%) and lower (3x)  $\bar{\beta}_{\text{inter-site}}$  and  $\bar{\beta}_{\text{inter-station}}$  (electronic supplementary material, table S5 and figure S9).

Beyond the hierarchical partitioning of diversity, we compared the distribution of fish MOTUs and species visual occurrences independently of the survey method and sampling effort using global species abundance distributions (gSAD) [37]. We fitted the fish MOTU and species visual occurrences to three distributions (log-series, Pareto and Pareto with exponential finite adjustment, i.e. Pareto Bended; see Methods) and estimated the parameters by maximum likelihood. For the visual census gSAD, the best

fit was obtained with the log-series and Pareto distributions (electronic supplementary material, table S6) with a slope of  $-0.95$  (confidence interval at 95%  $[-0.98; -0.92]$ ) (electronic supplementary material, figure S10). This suggests a distribution of geographically restricted or rare species close to the neutral theory ( $\beta$  close to  $-1$ ). By contrast, the best fit for fish MOTUs was obtained with the Pareto Bended distribution with a slope  $\beta = -0.76$  (confidence interval at 95%  $[-0.85; -0.65]$ ) and then with the log-series distribution, suggesting a lower prevalence of rarity than under the neutral theory, in agreement with previous tests based on species distributions on coral reefs [38].

### 3. Discussion

Environmental DNA allows the detection and identification of more taxa than traditional techniques [26,39], but further offers novel insights on the spatial organization of the richest marine ecosystem at a large scale. Over a timespan of 2.3 years, in major tropical ocean basins, eDNA metabarcoding reveals a higher proportion of crypto-benthic, pelagic and soft-sediment-associated fishes on coral reefs than detected in the most extensive visual census over 13 years. We found a high local MOTU turnover, but we were not able to conclude if it is due to an insufficient sampling at the station level, or if it suggests that differences in fish species composition may exist between adjacent reefs that are not detected



divers or are not permanent residents on coral reefs so can be missed in visual surveys [40]. Some crypto-benthic or reef-associated species, hidden in the reef, can also be missed by divers so were also more represented in eDNA than in visual surveys. Crypto-benthic species also have a crucial role for coral reef functioning, by promoting biomass production and fuelling the reef trophodynamics [41], but their diversity has been underestimated so far [13]. Transient, pelagic and deep-water species may be very important for reef functioning, through pelagic larval stages or nocturnal migration up the reef slope [12,42,43], but their presence and role need further investigation. By contrast, visual census also detected many families not detected, or not identified, by eDNA, such as Acanthuridae, Blenniidae, Caesionidae, Chaenopsidae, Chaetodontidae, Labrisomidae, Labridae or Microdesmidae. This limited identification by eDNA can be due to the very low representation of these families in 12S reference databases (between 0 and 12%), or to the low resolution of the teleo marker for species of these families, so several species can share the same sequence and be grouped under the same MOTU. Environmental DNA may also be inappropriate to detect these species in the environment.

The finding of a strong regional effect on both species composition (figure 3) and species differentiation (figure 4) at a large scale is in agreement with visual surveys and previous knowledge [44], while the suggestion of a strong turnover at the local scale may be an unexpected result for coral reef fishes. This predominant role of large-scale bioregional differentiation explains the exceptional fish diversity on coral reefs, probably associated with long-term geological isolation [2]. Overall, the Tropical Northwestern Atlantic region has a very distinct MOTU composition compared to the four other regions (figure 3) with only 1.2% of MOTUs being shared between the Tropical Northwestern Atlantic and any other region, while 20% of MOTUs are shared between at least two Indo-Pacific regions (figure 4a). The isolation of the Tropical Northwestern Atlantic region can be explained by the hard vicariant barrier of the Isthmus of Panama [14,45], and a limited suitable area for coral reefs during the past quaternary glaciation. By contrast, the Indo-Pacific maintained extensive coral reef refuges that have served as centres of survival during ice-age periods [9].

The greater local compositional dissimilarity of reef fishes among adjacent stations with eDNA than with visual census may correspond to local environmental or habitat differences, to stochastic or random processes [46], or may be due to an insufficient sampling at the station level (electronic supplementary material, analyses, figure S6). A higher number of replicates per station would be necessary to characterize exhaustively the diversity at the station level and more confidently conclude on the local turnover hypothesis.

While our results confirm the potential of eDNA to monitor biodiversity in marine ecosystems, some limitations should be addressed in the future to fully exploit this potential. Completing public reference databases would improve the accuracy of taxonomic assignment, which is essential for a better estimation of biodiversity patterns. At such a large spatial scale, reference databases are far from exhaustive with only up to 13% of fish species sequenced on our marker [47], preventing assignment to the species level for 81% of our eDNA sequences. Using multiple markers is an alternative to the database limitation [48,49], but it is much more expensive. For these reasons, we used MOTUs curated by a combination

of a clustering algorithm and conservative abundance-based post-clustering filters. While uncurated MOTUs are prone to overestimate real diversity [50] and a given MOTU can represent several species within one cluster or several MOTUs belonging to one species, MOTUs with conservative curation have been shown to reflect the true level of fish diversity across scales in streams [51,52]. Additionally, some species share the same barcode sequence due to insufficient genetic differentiation on such a small mitochondrial marker [49]. This lack of taxonomic resolution combined with a conservative curated MOTUs pipeline can underestimate MOTUs richness. Moreover, some crypto-benthic or rare fish families are still underrepresented in public databases, and their diversity is potentially underestimated with eDNA (i.e. Blenniidae, Gobiesocidae, Chaenopsidae and Aploactinidae).

Differences in sampling method and in sample size might influence the detected biodiversity with eDNA. The lower volume of water sampled in the Western Coral Triangle region (2 l per sample, so 4 l per station using point-sampling instead of 2 km transect with 30 l elsewhere), could underestimate fish biodiversity. However, previous studies show that MOTU accumulation curves based on this dataset were close to the total fish diversity reported in this region [32]. Furthermore,  $\beta$ -diversity between samples within stations in each region indicates that dissimilarity between samples is not greater in the Western Coral Triangle than in other regions (electronic supplementary material, figure S11). To account for differences in sample size and obtain a balanced design, we performed sensitivity analyses by rarefying our complete dataset to (i) four stations for all sites and (ii) four sites per region after removing the lowest sampled region (Southeast Polynesia) (electronic supplementary material, analyses, figures S1–S4). We obtained similar patterns even after subsampling stations or sites. However, our site-based and station-based accumulation curves do not reach plateaus suggesting that our sampling effort was not sufficient to exhaustively estimate fish biodiversity for each site (electronic supplementary material, analyses, figure S5) and station (electronic supplementary material, analyses, figure S6). Twenty-five replicates (so, 12 stations in case of field duplicates) could accurately estimate biodiversity regionally due to high local turnover [53]. A higher number of eDNA samples would be necessary here to reach MOTU accumulation per site and station.

The transport and degradation of eDNA can also impact species detection. As some evidence suggests that eDNA from pelagic fishes degrades slower than from inshore species [54], we cannot exclude that eDNA from pelagic and deep-water families (e.g. Myctophidae) might disperse sufficiently with sea currents such that species living close to reef habitats are detected. Environmental DNA transport could also explain the detection of some freshwater fish families (i.e. Centrarchidae, Osphronemidae or Channidae) in a few samples located near an estuary or in an enclosed bay with freshwater inputs.

Better understanding and anticipating the effects of multiple threats to the marine environment depends on the temporal and spatial extent of our monitoring capacity in the vast ocean. Environmental DNA is a powerful tool to investigate biodiversity patterns at large scale and monitor biodiversity, but still benefits from the combination with complementary approaches as visual methods for an exhaustive biodiversity survey across space and time to keep pace with ongoing changes.

## 4. Methods

### (a) Environmental DNA collection and sample processing

Environmental DNA seawater samples were collected between 2017 and 2019, following a hierarchical pattern. A total of 226 eDNA samples (filters) were collected in 100 stations (gathering of replicates at the same location) located in 26 sites (groups of stations separated by at least 35 km) distributed across five tropical regions (electronic supplementary material, figures S1 and S2). Three different sampling methods were used comprising a 2 km-long sampling transect of 30 l (surface or bottom depth) or point samples of 2 l (electronic supplementary material, table S7 and Methods S1), and between 12 and 64 samples were collected by region. Filtration was performed with polyethersulfone filters, 0.2 µm pore size. For each sampling campaign, a strict contamination control protocol was followed in both field and laboratory stages [39]. Negative field controls were performed in multiple sites, and revealed no contamination from the boat or samplers.

### (b) eDNA extraction, amplification and sequencing

DNA extraction was performed in a dedicated DNA laboratory (SPYGEN, [www.spygen.com](http://www.spygen.com)) equipped with positive air pressure, UV treatment and frequent air renewal. Decontamination procedures were conducted before and after all manipulations. Detailed protocols of DNA extraction, amplification and sequencing can be found in the electronic supplementary material, Method S2 and in [32,39]. A teleost-specific 12S mitochondrial rRNA primer pair (teleo, forward primer—ACACCGCCCGTCACTCT, reverse primer—CTTCCGGTACACTTACCATG [39]) was used for the amplification of metabarcode sequences. As we analysed our data using MOTUs as a proxy for species to overcome genetic database limitations, we chose to amplify only one marker. Teleo marker has been shown to be the most appropriate for fish, owing to its high interspecific variability, and its short size allowing us to detect rare and degraded DNA reliably [39,49,55,56]. Twelve DNA amplifications PCR per sample were performed.

### (c) Bioinformatic analysis

Following sequencing, reads were processed using clustering and post-clustering cleaning to remove errors and estimate the number of species using MOTUs [51]. First, reads were assembled using *VSEARCH* [57], then demultiplexed and trimmed using *CUTADAPT* [58] and clustering was performed using *SWARM* v.2 [59] with a minimum distance of 1 mismatch between clusters. Taxonomic assignment of MOTUs was carried out using the lower common ancestor (LCA) algorithm *ecotag* implemented in the *OBITOOLS* toolkit [60] and the European nucleotide archive as a reference database (release 143, March 2020). Details on the bioinformatics analysis can be found in the electronic supplementary material, Methods S3. Taxonomic assignments obtained from the LCA algorithm at the species level were accepted if the percentage of similarity with the reference sequence was 100%, at the genus level if the similarity was between 90 and 99%, and at the family level if the similarity was greater than 85% following previous studies [32,61]. If these criteria were not met, the MOTU was left unassigned. Only 21% of assigned MOTUs are assigned to the family level with a similarity between 85 and 90% (electronic supplementary material, table S8).

### (d) Visual census data

The visual census survey data used here is a subset (2047 transects, in 219 sites, electronic supplementary material, figure S1) of the complete visual census data (3027 transects) provided by the

RLS [31] and comprises all species observed on standardized 50 m surveys at sites in tropical biogeographic realms between 2006 and 2017 (electronic supplementary material, methods S4) [62]. We selected only the most recent survey for each station and only transects with more than five per cent of coral cover. Two different sampling protocols were adapted to detect both reef and crypto-benthic fishes.

### (e) Statistical analysis

More details on the statistical analysis are available in the electronic supplementary material, methods S5.

Accumulation curves were calculated for species per 500 m<sup>2</sup> transect, MOTUs per eDNA sample, and families per transect and sample. We used the functions ‘specaccum’ and ‘fitspecaccum’ from the R package ‘vegan’ which calculates the expected species accumulation curve using a sample-based rarefaction method and fit a nonlinear accumulation model. In order to assess the impact of the irregular sampling on the estimates measured with accumulation curves, we subset randomly half of the transects in the three most sampled regions in Australia and calculated again the accumulation curves for species and families (electronic supplementary material, figure S12). The results were unchanged.

Linear regression models were fitted between the number of MOTUs per family in the eDNA dataset and the number of species per family in the visual census dataset, after  $\log(x + 1)$  transformation (figure 1e).

Accumulation curves were also calculated by sub-setting MOTUs belonging to crypto-benthic orders, or to pelagic families, for both datasets (figure 2). The asymptote was calculated as described above.

We performed dbRDA on family proportions, with *region* and *site richness* as explanatory variables, using the function *capscale* from the *vegan* package. We subset the visual census to select only the 68 sites that fell into the five regions in common with the eDNA dataset. Total dbRDA provided the effects of each of the variables and their interaction. We then calculated partial dbRDA to measure the effect of the region while correcting for the effect of site richness (figure 3; electronic supplementary material, table S3).

We applied an additive partitioning framework [63] to separate the total MOTUs diversity at the global scale ( $\gamma$  global) into contributions at smaller scales from regions to local richness:  $\gamma_{\text{global}} = \beta_{\text{inter-region}} + \text{mean } \beta_{\text{inter-site}} + \text{mean } \beta_{\text{inter-station}} + \text{mean } \bar{\alpha}_{\text{station}}$ . In this additive framework, the three levels of biodiversity [64] (i.e.  $\alpha$ ,  $\beta$  and  $\gamma$ ) are expressed with the same unit and consequently the contribution of  $\alpha$  and  $\beta$  diversity to total diversity ( $\gamma$ ) can be directly compared [65].

We analysed the distribution of fish MOTU and species occurrences using gSAD which plots, on a log–log scale, the number of species as a function of the number of observations [37].

**Data accessibility.** All eDNA data (except New Caledonia) and visual census data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.3xsj3txj2>. New Caledonia eDNA data are available in Zenodo, <https://doi.org/10.5281/zenodo.6381130>. Code used for the analyses is available at [https://github.com/virginieamarques/Global\\_eDNA](https://github.com/virginieamarques/Global_eDNA). The bioinformatic pipeline used to analyse the metabarcoding data has been published in [51].

**Authors' contributions.** L.M.: conceptualization, data curation, formal analysis, methodology, visualization, writing—original draft and writing—review and editing; V.M.: conceptualization, data curation, formal analysis, methodology, visualization, writing—original draft and writing—review and editing; D.M.: conceptualization, funding acquisition, project administration, resources, supervision, validation, writing—original draft and writing—review and editing; C.A.: resources and writing—review and editing; M.A.: resources and writing—review and editing; F.B.: resources and writing—review and editing; G.H.B.-P.: resources and writing—review and editing; T.D.:



data curation, formal analysis and writing—review and editing; G.J.E.: resources and writing—review and editing; J.G.: resources and writing—review and editing; P.-E.G.: data curation, methodology, software and writing—review and editing; R.H.: funding acquisition, resources and writing—review and editing; J.-B.J.: resources and writing—review and editing; K.: resources and writing—review and editing; E.M.: resources and writing—review and editing; G.M.: resources and writing—review and editing; M.M.: resources and writing—review and editing; A.P.F.: resources and writing—review and editing; L.P.: resources and writing—review and editing; R.D.S.-S.: resources and writing—review and editing; H.Y.S.: resources and writing—review and editing; A.V.: data curation, methodology, resources and writing—review and editing; L.V.: resources and writing—review and editing; I.B.V.: resources and writing—review and editing; L.P.: conceptualization, funding acquisition, investigation, project administration, supervision, validation, writing—original draft and writing—review and editing; S.M.: conceptualization, funding acquisition, investigation, project administration, supervision, validation, writing—original draft and writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

**Conflict of interest declaration.** All authors declare that there is no conflict of interest regarding the publication of this article.

## References

- Parravicini V *et al.* 2013 Global patterns and predictors of tropical reef fish species richness. *Ecography (Cop)* **36**, 1254–1262. (doi:10.1111/j.1600-0587.2013.00291.x)
- Cowman PF, Bellwood DR. 2013 The historical biogeography of coral reef fishes: global patterns of origination and dispersal. *J. Biogeogr.* **40**, 209–224. (doi:10.1111/jbi.12003)
- Cinner JE *et al.* 2020 Meeting fisheries, ecosystem function, and biodiversity goals in a human-dominated world. *Science* **368**, 307–311. (doi:10.1126/science.aax9412)
- Hoegh-Guldberg O *et al.* 2019 The human imperative of stabilizing global climate change at 1.5°C. *Science* **365**, eaaw6974. (doi:10.1126/science.aaz4390)
- Victor BC. 2015 How many coral reef fish species are there? Cryptic diversity and the new molecular taxonomy. In *Ecology of fishes on coral reefs*, pp. 76–88. Cambridge, UK: Cambridge University Press.
- Siqueira AC, Morais RA, Bellwood DR, Cowman PF. 2020 Trophic innovations fuel reef fish diversification. *Nat. Commun.* **11**, 1–11. (doi:10.1038/s41467-020-16498-w)
- Bellwood D, Wainwright P. 2002 The history and biogeography of fishes on coral reefs. In *Coral reef fishes: dynamics and diversity in a complex ecosystem* (ed. P Sale). San Diego, CA: Academic Press.
- Bellwood DR, Hughes TP. 2001 Regional-scale assembly rules and biodiversity of coral reefs. *Science* **292**, 1532–1534. (doi:10.1126/science.1058635)
- Pellissier L *et al.* 2014 Quaternary coral reef refugia preserved fish diversity. *Science* **344**, 1016–1020. (doi:10.1126/science.1249853)
- Veron J, Devantier LM, Turak E, Green AL, Kininmonth S, Stafford-Smith M, Peterson N. 2009 Delineating the coral triangle. *Galaxea. J. Coral Reef Stud.* **11**, 91–100. (doi:10.3755/galaxea.11.91)
- Siqueira AC, Morais RA, Bellwood DR, Cowman PF. 2021 Planktivores as trophic drivers of global coral reef fish diversity patterns. *Proc. Natl Acad. Sci. USA* **118**, e2019404118. (doi:10.1073/pnas.2019404118)
- Morais RA, Bellwood DR. 2019 Pelagic subsidies underpin fish productivity on a degraded coral reef. *Curr. Biol.* **29**, 1521–1527. (doi:10.1016/j.cub.2019.03.044)
- Brandl SJ, Goatley CHR, Bellwood DR, Tornabene L. 2018 The hidden half: ecology and evolution of cryptobenthic fishes on coral reefs. *Biol. Rev.* **93**, 1846–1873. (doi:10.1111/brv.12423)
- Bender MG, Leprieur F, Mouillot D, Kulbicki M, Parravicini V, Pie MR, Barneche DR, Oliveira-Santos LGR, Floeter SR. 2017 Isolation drives taxonomic and functional nestedness in tropical reef fish faunas. *Ecography (Cop)* **40**, 425–435. (doi:10.1111/ecog.02293)
- Hughes TP, Bellwood DR, Connolly SR, Cornell HV, Karlson RH. 2014 Double jeopardy and global extinction risk in corals and reef fishes. *Curr. Biol.* **24**, 2946–2951. (doi:10.1016/j.cub.2014.10.037)
- Mouillot D, Villéger S, Parravicini V, Kulbicki M, Arias-gonzález JE. 2014 Functional over-redundancy and high functional vulnerability in global fish faunas on tropical reefs. *Proc. Natl Acad. Sci. USA* **111**, 13 757–13 762. (doi:10.1073/pnas.1317625111)
- Alzate A, Zapata FA, Giraldo A. 2014 A comparison of visual and collection-based methods for assessing community structure of coral reef fishes in the Tropical Eastern Pacific. *Rev. Biol. Trop.* **62**, 359–371. (doi:10.15517/rbt.v62i0.16361)
- Bellwood D, Renema W, Rosen BB. 2012 Biodiversity hotspots, evolution and coral reef biogeography: a review. In *Biotic evolution and environmental change in Southeast Asia*, pp. 216–245. Cambridge, UK: Cambridge University Press.
- Mora C. 2015 Ecology of fishes on coral reefs. In *Ecology of fishes on coral reefs*, pp. 1–374. Cambridge, UK: Cambridge University Press.
- Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH. 2012 Environmental DNA. *Mol. Ecol.* **21**, 1789–1793. (doi:10.1111/j.1365-294X.2012.05542.x)
- Boulanger E *et al.* 2021 Environmental DNA metabarcoding reveals and unpacks a biodiversity conservation paradox in Mediterranean marine reserves [Dryad Dataset]. *Proc. R. Soc. B* **288**, 20210112. (doi:10.1098/rsob.2021.0112)
- Harrison JB, Sunday JM, Rogers SM. 2019 Predicting the fate of eDNA in the environment and implications for studying biodiversity. *Proc. R. Soc. B* **286**, 1–9. (doi:10.1098/rsob.2019.1409)
- Cordier T *et al.* 2020 Ecosystem monitoring powered by environmental genomics: a review of current strategies with an implementation roadmap. *Mol. Biol. Evol.* **30**, 2937–2958. (doi:10.1111/mec.15472 1–22)
- De Vargas C *et al.* 2015 Eukaryotic plankton diversity in the sunlit ocean. *Science* **348**, 1–11. (doi:10.1126/science.1261605)
- Valdivia-Carrillo T, Rocha-Olivares A, Reyes-Bonilla H, Domínguez-Contreras JF, Munguia-Vega A. 2021 Integrating eDNA metabarcoding and simultaneous underwater visual surveys to describe complex fish communities in a marine biodiversity hotspot. *Mol. Ecol. Resour.* **21**, 1558–1574. (doi:10.1111/1755-0998.13375)

26. West K *et al.* 2021 Large-scale eDNA metabarcoding survey reveals marine biogeographic break and transitions over tropical north-western Australia. *Divers. Distrib.* **27**, 1942–1957. (doi:10.1111/ddi.13228)
27. Kume M *et al.* 2021 Factors structuring estuarine and coastal fish communities across Japan using environmental DNA metabarcoding. *Ecol. Indic.* **121**, 107216. (doi:10.1016/j.ecolind.2020.107216)
28. Fraija-Fernández N, Bouquieaux MC, Rey A, Mendibil I, Cotano U, Irigoien X, Santos M, Rodríguez-Ezpeleta N. 2020 Marine water environmental DNA metabarcoding provides a comprehensive fish diversity assessment and reveals spatial patterns in a large oceanic area. *Ecol. Evol.* **10**, 7560–7584. (doi:10.1002/ece3.6482)
29. Aglieri G *et al.* 2020 Environmental DNA effectively captures functional diversity of coastal fish communities. *Mol. Ecol.* **30**, 3127–3139. (doi:10.1111/mec.15661 1–13)
30. DiBattista JD *et al.* 2021 Environmental DNA reveals a multi-taxa biogeographic break across the Arabian Sea and Sea of Oman. *Environ. DNA* **4**, 206–221. (doi:10.1002/edn3.252)
31. Edgar GJ, Stuart-Smith RD. 2014 Systematic global assessment of reef fish communities by the Reef Life Survey program. *Sci. Data* **1**, 1–8. (doi:10.1038/sdata.2014.7)
32. Juhel JB *et al.* 2020 Accumulation curves of environmental DNA sequences predict coastal fish diversity in the coral triangle. *Proc. Biol. Sci.* **287**, 1–10. (doi:10.22541/au.158082947.75232617)
33. Castellanos-Galindo GA, Krumme U, Rubio EA, Saint-Paul U. 2013 Spatial variability of mangrove fish assemblage composition in the tropical eastern Pacific Ocean. *Rev. Fish Biol. Fish.* **23**, 69–86. (doi:10.1007/s11160-012-9276-4)
34. Willis TJ, Anderson MJ. 2003 Structure of cryptic reef fish assemblages: relationships with habitat characteristics and predator density. *Mar. Ecol. Prog. Ser.* **257**, 209–221. (doi:10.3354/meps257209)
35. Harnik PG, Simpson C, Payne JL. 2012 Long-term differences in extinction risk among the seven forms of rarity. *Proc. R. Soc. B* **279**, 4969–4976. (doi:10.1098/rspb.2012.1902)
36. Crist TO, Veech JA. 2006 Additive partitioning of rarefaction curves and species-area relationships: unifying  $\alpha$ -,  $\beta$ - and  $\gamma$ -diversity with sample size and habitat area. *Ecol. Lett.* **9**, 923–932. (doi:10.1111/j.1461-0248.2006.00941.x)
37. Enquist BJ *et al.* 2019 The commonness of rarity: global and future distribution of rarity across land plants. *Sci. Adv.* **5**, 1–14. (doi:10.1126/sciadv.aaz0414)
38. Dornelas M, Connolly SR, Hughes TP. 2006 Coral reef diversity refutes the neutral theory of biodiversity. *Nature* **440**, 80–82. (doi:10.1038/nature04534)
39. Valentini A *et al.* 2016 Next-generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. *Mol. Ecol.* **25**, 929–942. (doi:10.1111/mec.13428)
40. Boussarie G *et al.* 2018 Environmental DNA illuminates the dark diversity of sharks. *Sci. Adv.* **4**, 1–8. (doi:10.1126/sciadv.aap9661)
41. Brandl SJ *et al.* 2019 Demographic dynamics of the smallest marine vertebrates fuel coral-reef ecosystem functioning. *Science*, **364**(6446), 1189–1192.
42. Kimmerling N *et al.* 2018 Quantitative species-level ecology of reef fish larvae via metabarcoding. *Nat. Ecol. Evol.* **2**, 306–316. (doi:10.1038/s41559-017-0413-2)
43. Beckley LE, Holliday D, Sutton AL, Weller E, Olivier MP, Thompson PA. 2019 Structuring of larval fish assemblages along a coastal-oceanic gradient in the macro-tidal, tropical Eastern Indian Ocean. *Deep Res. Part II* **161**, 105–119. (doi:10.1016/j.dsr2.2018.03.008)
44. McLean M, Stuart-Smith RD, Villéger S, Auber A, Edgar GJ, MacNeil MA, Loiseau N, Leprieur F, Mouillot D. 2021 Trait similarity in reef fish faunas across the world's oceans. *Proc. Natl Acad. Sci. USA* **118**, e2012318118. (doi:10.1073/pnas.2012318118)
45. Gaboriau T, Leprieur F, Mouillot D, Hubert N. 2018 Influence of the geography of speciation on current patterns of coral reef fish biodiversity across the Indo-Pacific. *Ecography (Cop)* **41**, 1295–1306. (doi:10.1111/ecog.02589)
46. Ahmadi GN, Tornabene L, Smith DJ, Pezold FL. 2018 The relative importance of regional, local, and evolutionary factors structuring cryptobenthic coral-reef assemblages. *Coral Reefs* **37**, 279–293. (doi:10.1007/s00338-018-1657-2)
47. Marques V, Milhau T, Albouy C, Dejean T, Manel S, Mouillot D, Juhel JB. 2021 GApE DNA: assessing and mapping global species gaps in genetic databases for eDNA metabarcoding. *Divers. Distrib.* **27**, 1880–1892. (doi:10.1111/ddi.13142)
48. Ruppert KM, Kline RJ, Rahman MS. 2019 Past, present, and future perspectives of environmental DNA (eDNA) metabarcoding: a systematic review in methods, monitoring, and applications of global eDNA. *Glob. Ecol. Conserv.* **17**, e00547. (doi:10.1016/j.gecco.2019.e00547)
49. Polanco A *et al.* 2021 Comparing the performance of 12S mitochondrial primers for fish environmental DNA across ecosystems. *Environ. DNA* **3**, 1113–1127. (doi:10.1002/edn3.232)
50. Brandt MI, Trouche B, Quintric L, Günther B, Wincker P, Poulain J, Arnaud-Haond S. 2021 Bioinformatic pipelines combining denoising and clustering tools allow for more comprehensive prokaryotic and eukaryotic metabarcoding. *Mol. Ecol. Resour.* **21**, 1904–1921. (doi:10.1111/1755-0998.13398)
51. Marques V, Guérin PÉ, Rocle M, Valentini A, Manel S, Mouillot D, Dejean T. 2020 Blind assessment of vertebrate taxonomic diversity across spatial scales by clustering environmental DNA metabarcoding sequences. *Ecography (Cop)* **43**, 1779–1790. (doi:10.1111/ecog.05049)
52. Sales NG, Wangenstein OS, Carvalho DC, Deiner K, Praebel I, McDevitt A, Mariani S. 2020 Space-time dynamics in monitoring neotropical fish communities using eDNA metabarcoding. *bioRxiv*.
53. Stauffer S *et al.* 2021 How many replicates to accurately estimate fish biodiversity using environmental DNA on coral reefs? *bioRxiv*.
54. Collins RA, Wangenstein OS, O'Gorman EJ, Mariani S, Sims DW, Genner MJ. 2018 Persistence of environmental DNA in marine systems. *Commun. Biol.* **1**, 1–12. (doi:10.1038/s42003-018-0192-6)
55. Collins RA, Bakker J, Wangenstein OS, Soto AZ, Corrigan L, Sims DW, Genner MJ, Mariani S. 2019 Non-specific amplification compromises environmental DNA metabarcoding with COI. *Methods Ecol. Evol.* **10**, 1985–2001. (doi:10.1111/2041-210X.13276)
56. Zhang S, Zhao J, Yao M. 2020 A comprehensive and comparative evaluation of primers for metabarcoding eDNA from fish. *Methods Ecol. Evol.* **2020**, 1609–1625. (doi:10.1111/2041-210X.13485)
57. Rognes T, Flourie T, Nichols B, Quince C, Mahé F. 2016 VSEARCH: a versatile open source tool for metagenomics. *PeerJ* **4**, 1–22. (doi:10.7717/peerj.2584)
58. Martin M. 1994 Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet.journal* **17**, 10–12. (doi:10.14806/ej.17.1.200)
59. Mahé F, Rognes T, Quince C, de Vargas C, Dunthorn M. 2015 Swarm v2: highly-scalable and high-resolution amplicon clustering. *PeerJ* **3**, 1–12. (doi:10.7717/peerj.1420)
60. Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P. 2016 Coissac E. obitools: a unix-inspired software package for DNA metabarcoding. *Mol. Ecol. Resour.* **16**, 176–182. (doi:10.1111/1755-0998.12428)
61. Polanco Fernández A *et al.* 2020 Comparing environmental DNA metabarcoding and underwater visual census to monitor tropical reef fishes. *Environ. DNA* **3**, 142–156. (doi:10.1002/edn3.140)
62. Spalding MD *et al.* 2007 Marine ecoregions of the World: a bioregionalization of coastal and shelf areas. *Bioscience* **57**, 573–583. (doi:10.1641/B570707)
63. Escalas A *et al.* 2017 Functional diversity and redundancy across fish gut, sediment and water bacterial communities. *Environ. Microbiol.* **19**, 3268–3282. (doi:10.1111/1462-2920.13822)
64. Whittaker RH. 1972 Evolution and measurement of species diversity. *Taxon* **21**, 213–251. (doi:10.2307/1218190)
65. Veech JA, Summerville KS, Crist TO, Gering JC. 2002 The additive partitioning of species diversity: recent revival of an old idea. *Oikos* **99**, 3–9. (doi:10.1034/j.1600-0706.2002.990101.x)